

# ***KillTest***

Higher Quality, Better Service!



## **Q&A**

<http://www.killtest.com>

We offer free update service for one year.

**Exam** : **IFoA\_CAA\_M0**

**Title** : **Module 0 - Entry Exam**

**Version** : **DEMO**

1 .1/5 of actuarial students like skiing. 2/5of actuarial students like snowboarding. 1/3 of actuarial students like skiing and snowboarding.

Calculate the proportion of actuarial students that like skiing or snowboarding.

A)

$$-\frac{4}{15}$$

B)

$$\frac{4}{15}$$

C)

$$\frac{3}{5}$$

D)

$$\frac{14}{15}$$

A. Option A

B. Option B

C. Option C

D. Option D

**Answer: B**

2 .Calculate the sum of the following non-terminating progression:

2/10, 2/40, 2/160, 2/640,...

A. 0.174

B. 0.266

C. 0.267

D. 0.406

**Answer: C**

3 .For random variable X, use the following statistics to calculate its coefficient of skewness based on central moments.

$$E(X) = 3.940$$

$$E(X^2) = 21.466$$

$$\text{skew}(X) = E[(X - )^3] = 6.008$$

A. -0.415

B. 0.060

C. 0.415

D. 0.768

**Answer: C**

4 .Identify the condition that fully describes the existence of independence between two events A and B.

- A.  $P(A|B) = P(A)/P(B)$  and  $P(B|A) = P(B)/P(A)$
- B.  $P(A|B) = P(A) - P(B)$  and  $P(B|A) = P(B) - P(A)$
- C.  $P(A|B) = P(A)$  and  $P(B|A) = P(B)$
- D.  $P(A|B) = P(A) + P(B)$  and  $P(B|A) = P(B) + P(A)$

**Answer: C**

5 .Determine which of the options is equal to  $\log(3) - 2\log(x+1)$ .

A)

$$\log(2x + 1)$$

B)

$$\log\left(\frac{3}{2x + 1}\right)$$

C)

$$\log\left(3(x + 1)^2\right)$$

D)

$$\log\left(\frac{3}{(x + 1)^2}\right)$$

- A. Option A
- B. Option B
- C. Option C
- D. Option D

**Answer: D**